

PART 1 - CHEMISTRY

Topic : Surface chemistry

SECTION-B : (Maximum Marks : 8)

This section contains **TWO** questions.

The answer to each question is a **NUMERICAL VALUE**.

For each question, enter the correct numerical value of the answer (in decimal notation, truncated / rounded-off to the **SECOND DECIMAL PLACE**; e.g. 6.25, 7.00, -0.33, -.30, 30.27, -127.30) in the place designated to enter the answer.

Answer to each question will be evaluated according to the following marking scheme :

Full Marks : +4 If **ONLY** the correct answer is entered;

Zero Marks : 0 In all other cases.

1. At a given temperature, 0.45 g of acetic acid in 50 mL of water is shaken with 1.0 g of charcoal and the pH of the resulting solution is 3.0. Assume, the adsorption of acetic acid from the aqueous solution by charcoal follows Freundlich isotherm,

$$\frac{x}{m} = kC^{1/n}$$

If the plot of $\log_{10}(x/m)$ against $\log_{10} C$ gives a straight line with slope 1, the value of k in L mol^{-1} is ____.

Given: The molar mass of acetic acid is 60 g mol^{-1} .

The acid dissociation constant of acetic acid is 1.0×10^{-5} at the given temperature.

x is the mass (in grams) of acetic acid adsorbed.

m is the mass (in grams) of charcoal.

C is the equilibrium concentration of acetic acid in the solution after the adsorption is complete.

k and n are constants for acetic acid–charcoal system at the given temperature.

[JEE Advanced 2026 Paper 2]

Scan or tap for the
full solution



2. Adsorption of phenol from its aqueous solution on to fly ash obeys Freundlich isotherm. At a given temperature, from 10 mg g^{-1} and 16 mg g^{-1} aqueous phenol solutions, the concentrations of adsorbed phenol are measured to be 4 mg g^{-1} and 10 mg g^{-1} , respectively. At this temperature, the concentration (in mg g^{-1}) of adsorbed phenol from 20 mg g^{-1} aqueous solution of phenol will be ____.

Use: $\log_{10} 2 = 0.3$

[JEE Advanced 2025 Paper 2]

Scan or tap for the
full solution



Space for Rough Work

**JEE ADVANCED SURFACE CHEMISTRY —
FREUNDLICH ADSORPTION ISOTHERM — PYQS**

Test Pattern : JEE (Advanced)

ANSWER KEY**SECTION-B**

Ques.	1	2								
Ans.	1.45 to 1.52	15.5 to 16.5								