

PYQSide

NEET PRINCIPLES OF INHERITANCE AND VARIATION — DIHYBRID CROSS AND LAW OF INDEPENDENT ASSORTMENT — PYQS

Max. Marks: 48

Questions: 12

Section A (MCQ)

Q.1 Genes R and Y follow independent assortment. If RRY Y produce round yellow seeds and rryy produce wrinkled green seeds, what will be the phenotypic ratio of the F₂ generation?

[NEET (UG) 2025]

- (1) Phenotypic ratio - 9:3:3:1
- (2) Phenotypic ratio - 9:7
- (3) Phenotypic ratio - 1:2:1
- (4) Phenotypic ratio - 3:1

Scan or tap for the full solution 

Q.2 When a tall pea plant with round seeds was selfed, it produced the progeny of :

- (a) Tall plants with round seeds and
- (b) Tall plants with wrinkled seeds.

Identify the genotype of the parent plant.

[RE-NEET (UG) 2024]

- (1) TtRr
- (2) TtRR
- (3) TTRR
- (4) TTRr

Scan or tap for the full solution 

Q.3 What is the expected percentage of F₂ progeny with yellow and inflated pod in dihybrid cross experiment involving pea plants with green coloured, inflated pod and yellow coloured constricted pod?

[NEET (UG) 2022 (Phase 2)]

- (1) 9%
- (2) 100%
- (3) 56.25%
- (4) 18.75%

Scan or tap for the full solution 

Q.4 A human male produces sperms with the genotypes AB, Ab, aB, and ab pertaining to two diallelic characters in equal proportions. What is the corresponding genotype of this person?

[AIPMT 2007]

- (1) AaBb
- (2) AABB
- (3) AaBB
- (4) AABb

Scan or tap for the full solution 

Q.5 Test cross involves-

[AIPMT 2006]

- (1) Crossing between two F₁ hybrids
- (2) Crossing the F₁ hybrid with a double recessive genotype
- (3) Crossing between two genotypes with recessive trait
- (4) Crossing between two genotypes with dominant trait

Scan or tap for the full solution 

Q.6 In Mendel's experiments with garden pea, round seed shape (RR) was dominant over wrinkled seeds (rr), yellow cotyledon (YY) was dominant over green cotyledon (yy). What are the expected phenotypes in the F₂ generation of the cross RRY Y × rryy?

[AIPMT 2006]

- (1) Round seeds with yellow cotyledons, and wrinkled seeds with yellow cotyledons
- (2) Only wrinkled seeds with green cotyledons
- (3) Only wrinkled seeds with yellow cotyledons
- (4) Only round seeds with green cotyledons

Scan or tap for the full solution 

Q.7 How many different kinds of gametes will be produced by a plant having the genotype AABbCC?

[AIPMT 2006]

- (1) Two
- (2) Nine
- (3) Four
- (4) Three

Scan or tap for the full solution 

Q.8 In order to find out the different types of gametes produced by a pea plant having the genotype AaBb, it should be crossed to a plant with the genotype -

[AIPMT 2005]

- (1) aaBB
- (2) aabb
- (3) AaBb
- (4) AABB

Scan or tap for the full solution 

Q.9 In a plant, red fruit (R) is dominant over yellow fruit (r) and tallness (T) is dominant over shortness (t). If a plant with RRTt genotype is crossed with a plant that is rrtt

[AIPMT 2004]

- (1) 75% will be tall with red fruit
- (2) 25% will be tall with red fruit
- (3) 50% will be tall with red fruit
- (4) All the offspring will be tall with red fruit

Scan or tap for the
full solution



Q.10 Two non-allelic genes produce the new phenotype when present together but fail to do so independently then it is called [AIPMT 2001]

- (1) Polygene
- (2) Complementary gene
- (3) Non-complementary gene
- (4) Epistasis

Scan or tap for the
full solution



Q.11 Ratio of complementary genes is [AIPMT 2001]

- (1) 9 : 3 : 4
- (2) 9 : 7
- (3) 12 : 3 : 1
- (4) 9 : 3 : 3 : 4

Scan or tap for the
full solution



Q.12 Due to the cross between $TTRr \times ttrr$ the resultant progenies showed how many percent plants tall, red flowered: [AIPMT 2000]

- (1) 50%
- (2) 100%
- (3) 75%
- (4) 25%

Scan or tap for the
full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9	10	11
Ans.	1	4	4	1	2	1	1	2	3	2	2
Ques.	12										
Ans.	1										