

PYQSide

NEET SEXUAL REPRODUCTION IN FLOWERING PLANTS — THE PISTIL, MEGASPORANGIUM (OVULE) AND EMBRYO SAC (MEGASPOROGENESIS) — PYQS

Max. Marks: 88

Questions: 22

Section A (MCQ)

Q.1 Which of the following in female gametophyte of an angiosperm helps in guiding the pollen tube for fertilizing the eggs? [RE-NEET (UG) 2026]

- (1) Polar nucleus (2) Antipodals
(3) Synergids (4) Central cells

Scan or tap for the full solution



Q.2 Given below are two statements : One is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

Assertion (A): A typical unfertilised, angiosperm embryo sac at maturity is 8 nucleate and 7-celled.

Reason (R): The egg apparatus has 2 polar nuclei.

In the light of the above statements, choose the correct answer from the options given below :

[NEET (UG) 2025]

- (1) A is true but R is false
(2) A is false but R is true
(3) Both A and R are true and R is the correct explanation of A
(4) Both A and R are true but R is NOT the correct explanation of A

Scan or tap for the full solution



Q.3 How many meiotic and mitotic divisions need to occur for the development of a mature female gametophyte from the megaspore mother cell in an angiosperm plant? [NEET (UG) 2025]

- (1) 1 Meiosis and 3 Mitosis
(2) No Meiosis and 2 Mitosis
(3) 2 Meiosis and 3 Mitosis
(4) 1 Meiosis and 2 Mitosis

Scan or tap for the full solution



Q.4 Which part of the ovule stores reserve food materials? [RE-NEET (UG) 2024]

- (1) Nucellus (2) Integument
(3) Placenta (4) Funicle

Scan or tap for the full solution



Q.5 In general the egg apparatus of embryo sac in angiosperm consists of [NEET (UG) 2022 (Phase 2)]

- (1) One egg cell, two synergids, two antipodal cells, two Polar nuclei
(2) One egg cell, two synergids, three antipodal cells, two Polar nuclei
(3) One egg cell, two synergids, two antipodal cells, three Polar nuclei
(4) One egg cell, three synergids, two antipodal cells, two Polar nuclei

Scan or tap for the full solution



Q.6 A typical angiosperm embryo sac at maturity is :

[NEET (UG) 2021]

- (1) 8-nucleate and 8-celled
(2) 8-nucleate and 7-celled
(3) 7-nucleate and 8-celled
(4) 7-nucleate and 7-celled

Scan or tap for the full solution



Q.7 The body of the ovule is fused with the funicle at

[NEET (UG) 2020]

- (1) Chalaza (2) Hilum
(3) Micropyle (4) Nucellus

Scan or tap for the full solution



Q.8 Functional megaspore in an angiosperm develops into an [NEET 2017]

- (1) endosperm (2) embryo sac
(3) embryo (4) ovule.

Scan or tap for the full solution



Q.9 In majority of angiosperms [NEET 2016 (Phase 2)]

- (1) egg has a filiform apparatus
(2) there are numerous antipodal cells
(3) reduction division occurs in the megaspore mother cells
(4) a small central cell is present in that embryo sac.

Scan or tap for the full solution



Q.10 The ovule of an angiosperm is technically equivalent to [NEET 2016 (Phase 2)]

- (1) megasporangium
- (2) megasporophyll
- (3) megaspore mother cell
- (4) megaspore.

Scan or tap for the full solution



Q.11 Filiform apparatus is characteristic feature of [AIPMT 2015]

- (1) aleurone cell
- (2) synergids
- (3) generative cell
- (4) nucellar embryo.

Scan or tap for the full solution



Q.12 Function of filiform apparatus is to [AIPMT 2014]

- (1) recognize the suitable pollen at stigma
- (2) stimulate division of generative cell
- (3) produce nectar
- (4) guide the entry of pollen tube.

Scan or tap for the full solution



Q.13 Megasporangium is equivalent to [NEET 2013]

- (1) nucellus
- (2) ovule
- (3) embryo sac
- (4) fruit.

Scan or tap for the full solution



Q.14 Megaspores are produced from the megaspore mother cells after [NEET 2013 (Karnataka)]

- (1) mitotic division
- (2) formation of thick wall
- (3) differentiation
- (4) meiotic division.

Scan or tap for the full solution



Q.15 In angiosperms, functional megaspore develops into [AIPMT 2011 (Mains)]

- (1) embryo sac
- (2) ovule
- (3) endosperm
- (4) pollen sac.

Scan or tap for the full solution



Q.16 Filiform apparatus is a characteristic feature of [AIPMT 2011 (Prelims)]

- (1) suspensor
- (2) egg
- (3) synergid
- (4) zygote.

Scan or tap for the full solution



Q.17 Which one of the following pairs of plant structures has haploid number of chromosomes? [AIPMT 2008]

- (1) Nucellus and antipodal cells
- (2) Egg nucleus and secondary nucleus
- (3) Megaspore mother cell and antipodal cells
- (4) Egg cell and antipodal cells

Scan or tap for the full solution



Q.18 The arrangement of the nuclei in a normal embryo sac in the dicot plants is [AIPMT 2006]

- (1) 3 + 3 + 2
- (2) 2 + 4 + 2
- (3) 3 + 2 + 3
- (4) 2 + 3 + 3

Scan or tap for the full solution



Q.19 Which one of the following represents an ovule, where the embryo sac becomes horse-shoe shaped and the funiculus and micropyle are close to each other? [AIPMT 2005]

- (1) Amphitropous
- (2) Circinotropous
- (3) Atropous
- (4) Anatropous

Scan or tap for the full solution



Q.20 An ovule which becomes curved so that the nucellus and embryo sac lie at right angles to the funicle is [AIPMT 2004]

- (1) hemitropous
- (2) campylotropous
- (3) anatropous
- (4) orthotropous.

Scan or tap for the full solution



Q.21 What is the direction of micropyle in anatropous ovule? [AIPMT 2002]

- (1) Upwards
- (2) Downward
- (3) Right
- (4) Left

Scan or tap for the full solution



Q.22 Eight nucleated embryo sac is [AIPMT 2000]

- (1) only monosporic
- (2) only bisporic
- (3) only tetrasporic
- (4) any of these.

Scan or tap for the full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9	10	11
Ans.	3	1	1	1	2	2	2	2	3	1	2
Ques.	12	13	14	15	16	17	18	19	20	21	22
Ans.	4	2	4	1	3	4	3	1	1	2	4