

# PYQSide

## NEET ATOMS — PYQS

Max. Marks: 196

Questions: 49

### Section A (MCQ)

- Q.1** In the first excited state of hydrogen atom, the energy of its electron is  $-3.4$  eV. The radial distance of the electron from the hydrogen nucleus in this case is approximately :

(Take  $1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$ ,  $e = 1.6 \times 10^{-19} \text{ C}$  and

$$\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ N m}^2/\text{C}^2 \quad [\text{NEET (UG) 2026}]$$

- (1)  $2.1 \times 10^{-8} \text{ m}$  (2)  $2.1 \times 10^{-10} \text{ m}$   
(3)  $2.1 \times 10^{-11} \text{ m}$  (4)  $2.1 \times 10^{-9} \text{ m}$

Scan or tap for the full solution



- Q.2** Consider that an electron is revolving in an excited state of Hydrogen atom with velocity

$\sqrt{25.6} \times 10^5 \text{ ms}^{-1}$ . The radius of the orbit is  $x \times 10^{-9} \text{ m}$ . The value of  $x$  is:

[Take the mass of electron to be  $9 \times 10^{-31} \text{ kg}$ , charge of electron =  $-1.6 \times 10^{-19} \text{ C}$  and





$$\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2/\text{C}^2 \quad [\text{RE-NEET (UG) 2026}]$$

- (1) 1 (2) 4 (3) 3 (4) 2

Scan or tap for the full solution



- Q.3** In Geiger-Marsden experiment, the number of scattered  $\alpha$ -particles  $N(\theta)$  is plotted as a function of scattering angle  $\theta$ . Which of the following options represents the correct plot? [RE-NEET (UG) 2026]



Scan or tap for the full solution



- Q.4** A particle of mass  $m$  is moving around the origin with a constant force  $F$  pulling it towards the origin. If Bohr model is used to describe its motion, the radius of the  $n^{\text{th}}$  orbit and the particle's speed  $v$  in the orbit depend on  $n$  as [NEET (UG) 2025]

- (1)  $r \propto n^{2/3}$ ;  $v \propto n^{1/3}$  (2)  $r \propto n^{4/3}$ ;  $v \propto n^{-1/3}$   
(3)  $r \propto n^{1/3}$ ;  $v \propto n^{1/3}$  (4)  $r \propto n^{1/3}$ ;  $v \propto n^{2/3}$

Scan or tap for the full solution



- Q.5** De-Broglie wavelength of an electron orbiting in the  $n = 2$  state of hydrogen atom is close to

(Given Bohr radius =  $0.052 \text{ nm}$ ) [NEET (UG) 2025]

- (1)  $1.67 \text{ nm}$  (2)  $2.67 \text{ nm}$   
(3)  $0.067 \text{ nm}$  (4)  $0.67 \text{ nm}$

Scan or tap for the full solution



- Q.6** Given below are two statements:

**Statement-I:** Atoms are electrically neutral as they contain equal number of positive and negative charges.

**Statement-II:** Atoms of each element are stable and emit their characteristic spectrum.

In the light of the above statements, choose the most appropriate answer from the options given below.

[NEET (UG) 2024]

- (1) Both Statement-I and Statement-II are correct  
(2) Both Statement-I and Statement-II are incorrect  
(3) Statement-I is correct but Statement-II is incorrect  
(4) Statement-I is incorrect but Statement-II is correct

Scan or tap for the full solution



- Q.7** Match List I with List II:

| List I<br>(Spectral Lines of Hydrogen<br>for transitions from) |                        | List II<br>(Wavelengths<br>(nm)) |       |
|----------------------------------------------------------------|------------------------|----------------------------------|-------|
| A.                                                             | $n_2 = 3$ to $n_1 = 2$ | I.                               | 410.2 |
| B.                                                             | $n_2 = 4$ to $n_1 = 2$ | II.                              | 434.1 |
| C.                                                             | $n_2 = 5$ to $n_1 = 2$ | III.                             | 656.3 |
| D.                                                             | $n_2 = 6$ to $n_1 = 2$ | IV.                              | 486.1 |

Choose the correct answer from the options given below: [NEET (UG) 2024]

- (1) A-II, B-I, C-IV, D-III (2) A-III, B-IV, C-II, D-I  
(3) A-IV, B-III, C-I, D-II (4) A-I, B-II, C-III, D-IV


Scan or tap for the full solution 

**Q.8** The spectral series which corresponds to the electronic transition from the levels  $n_2 = 5, 6, \dots$  to the level  $n_1 = 4$  is [RE-NEET (UG) 2024]

- (1) Pfund series (2) Brackett series  
(3) Lyman series (4) Balmer series

Scan or tap for the full solution 

**Q.9** Some energy levels of a molecule are shown in the figure with their wavelengths of transitions. Then : [RE-NEET (UG) 2024]



- (1)  $\lambda_3 > \lambda_2, \lambda_1 = 2\lambda_2$  (2)  $\lambda_3 > \lambda_2, \lambda_1 = 4\lambda_2$   
(3)  $\lambda_1 > \lambda_2, \lambda_2 = 2\lambda_3$  (4)  $\lambda_2 > \lambda_1, \lambda_2 = 2\lambda_3$

Scan or tap for the full solution 

**Q.10** In hydrogen spectrum, the shortest wavelength in the Balmer series is  $\lambda$ . The shortest wavelength in the Brackett series is : [NEET (UG) 2023]

- (1)  $4\lambda$  (2)  $9\lambda$  (3)  $16\lambda$  (4)  $2\lambda$

Scan or tap for the full solution 

**Q.11** The radius of inner most orbit of hydrogen atom is  $5.3 \times 10^{-11}$  m. What is the radius of third allowed orbit of hydrogen atom? [NEET (UG) 2023]

- (1)  $1.06 \text{ \AA}$  (2)  $1.59 \text{ \AA}$  (3)  $4.77 \text{ \AA}$  (4)  $0.53 \text{ \AA}$

Scan or tap for the full solution 

**Q.12** Let  $T_1$  and  $T_2$  be the energy of an electron in the first and second excited states of hydrogen atoms, respectively. According to the Bohr's model of an atom, the ratio  $T_1:T_2$  is [NEET (UG) 2022]

- (1) 1:4 (2) 4:1 (3) 4:9 (4) 9:4

Scan or tap for the full solution 

**Q.13** Let  $R_1$  be the radius of the second stationary orbit and  $R_2$  be the radius of the fourth stationary orbit of an electron in Bohr's model. The ratio  $\frac{R_1}{R_2}$  is : [NEET (UG) 2022 (Phase 2)]

- (1) 4 (2) 0.25 (3) 0.5 (4) 2

Scan or tap for the full solution 

**Q.14** For which one of the following, Bohr model is not valid? [NEET (UG) 2020]

- (1) Singly ionised helium atom ( $\text{He}^+$ )  
(2) Deuteron atom  
(3) Singly ionised neon atom ( $\text{Ne}^+$ )  
(4) Hydrogen atom

Scan or tap for the full solution 

**Q.15**  $\alpha$ -particle consists of : [NEET (UG) 2019]

- (1) 2 electrons, 2 protons and 2 neutrons  
(2) 2 electrons and 4 protons only  
(3) 2 protons only  
(4) 2 protons and 2 neutrons only

Scan or tap for the full solution 

**Q.16** The total energy of an electron in an atom in an orbit is  $-3.4$  eV. Its kinetic and potential energies are, respectively. [NEET (UG) 2019]

- (1) 3.4 eV,  $-6.8$  eV (2) 3.4 eV, 3.4 eV  
(3)  $-3.4$  eV,  $-3.4$  eV (4)  $-3.4$  eV,  $-6.8$  eV

Scan or tap for the full solution 

**Q.17** The ratio of kinetic energy to the total energy of an electron in a Bohr orbit of the hydrogen atom, is [NEET 2018]

- (1) 1:1 (2) 1: -1 (3) 2: -1 (4) 1: -2

Scan or tap for the full solution 

**Q.18** The ratio of wavelengths of the last line of Balmer series and the last line of Lyman series is [NEET 2017]

- (1) 1 (2) 4 (3) 0.5 (4) 2

Scan or tap for the full solution 

**Q.19** Given the value of Rydberg constant is  $10^7 \text{ m}^{-1}$ , the wave number of the last line of the Balmer series in hydrogen spectrum will be [NEET 2016 (Phase 1)]

- (1)  $0.25 \times 10^7 \text{ m}^{-1}$  (2)  $2.5 \times 10^7 \text{ m}^{-1}$   
(3)  $0.025 \times 10^4 \text{ m}^{-1}$  (4)  $0.5 \times 10^7 \text{ m}^{-1}$

Scan or tap for the full solution 

**Q.20** When an  $\alpha$ -particle of mass  $m$  moving with velocity  $v$  bombards on a heavy nucleus of charge  $Ze$ , its distance of closest approach from the nucleus depends on  $m$  as [NEET 2016 (Phase 1)]

- (1)  $\frac{1}{m^2}$       (2)  $m$       (3)  $\frac{1}{m}$       (4)  $\frac{1}{\sqrt{m}}$

Scan or tap for the full solution



**Q.21** Electrons of mass  $m$  with de-Broglie wavelength  $\lambda$  fall on the target in an X-ray tube. The cutoff wavelength ( $\lambda_0$ ) of the emitted X-ray is

[NEET 2016 (Phase 2)]

- (1)  $\lambda_0 = \frac{2mc\lambda^2}{h}$       (2)  $\lambda_0 = \frac{2h}{mc}$   
 (3)  $\lambda_0 = \frac{2m^2c^2\lambda^3}{h^2}$       (4)  $\lambda_0 = \lambda$

Scan or tap for the full solution



**Q.22** If an electron in a hydrogen atom jumps from the 3<sup>rd</sup> orbit to the 2<sup>nd</sup> orbit, it emits a photon of wavelength  $\lambda$ . When it jumps from the 4<sup>th</sup> orbit to the 3<sup>rd</sup> orbit, the corresponding wavelength of the photon will be

[NEET 2016 (Phase 2)]

- (1)  $\frac{16}{25}\lambda$       (2)  $\frac{9}{16}\lambda$       (3)  $\frac{20}{7}\lambda$       (4)  $\frac{20}{13}\lambda$

Scan or tap for the full solution



**Q.23** In the spectrum of hydrogen, the ratio of the longest wavelength in the Lyman series to the longest wavelength in the Balmer series is

[AIPMT 2015]

- (1)  $\frac{27}{5}$       (2)  $\frac{5}{27}$       (3)  $\frac{4}{9}$       (4)  $\frac{9}{4}$

Scan or tap for the full solution



**Q.24** Consider 3<sup>rd</sup> orbit of  $\text{He}^+$  (Helium), using non-relativistic approach, the speed of electron in this orbit will be [given  $K = 9 \times 10^9$  constant,  $Z = 2$  and  $h$  (Planck's Constant)  $= 6.6 \times 10^{-34}$  J s]

[AIPMT 2015 (Cancelled)]

- (1)  $0.73 \times 10^6$  m/s      (2)  $3.0 \times 10^8$  m/s  
 (3)  $2.92 \times 10^6$  m/s      (4)  $1.46 \times 10^6$  m/s

Scan or tap for the full solution



**Q.25** Hydrogen atom in ground state is excited by a monochromatic radiation of  $\lambda = 975 \text{ \AA}$ . Number of spectral lines in the resulting spectrum emitted will be

[AIPMT 2014]

- (1) 3      (2) 2      (3) 6      (4) 10

Scan or tap for the full solution



**Q.26** Ratio of longest wave lengths corresponding to Lyman and Balmer series in hydrogen spectrum is

[NEET 2013]

- (1)  $\frac{7}{29}$       (2)  $\frac{9}{31}$       (3)  $\frac{5}{27}$       (4)  $\frac{3}{23}$

Scan or tap for the full solution



**Q.27** An electron in hydrogen atom makes a transition  $n_1 \rightarrow n_2$  where  $n_1$  and  $n_2$  are principal quantum numbers of the two states. Assuming Bohr's model to be valid, the time period of the electron in the initial state is eight times that in the final state. The possible values of  $n_1$  and  $n_2$  are

[NEET 2013 (Karnataka)]

- (1)  $n_1 = 6$  and  $n_2 = 2$       (2)  $n_1 = 8$  and  $n_2 = 1$   
 (3)  $n_1 = 8$  and  $n_2 = 2$       (4)  $n_1 = 4$  and  $n_2 = 2$

Scan or tap for the full solution



**Q.28** The transition from the state  $n = 3$  to  $n = 1$  in a hydrogen like atom results in ultraviolet radiation. Infrared radiation will be obtained in the transition from

[AIPMT 2012 (Mains)]

- (1)  $2 \rightarrow 1$       (2)  $3 \rightarrow 2$   
 (3)  $4 \rightarrow 2$       (4)  $4 \rightarrow 3$

Scan or tap for the full solution



**Q.29** Electron in hydrogen atom first jumps from third excited state to second excited state and then from second excited to the first excited state. The ratio of the wavelengths  $\lambda_1 : \lambda_2$  emitted in the two cases is

[AIPMT 2012 (Prelims)]

- (1)  $\frac{7}{5}$       (2)  $\frac{27}{20}$       (3)  $\frac{27}{5}$       (4)  $\frac{20}{7}$

Scan or tap for the full solution



**Q.30** An electron of a stationary hydrogen atom passes from the fifth energy level to the ground level. The velocity that the atom acquired as a result of photon emission will be

[AIPMT 2012 (Prelims)]

- (1)  $\frac{24hR}{25m}$       (2)  $\frac{25hR}{24m}$       (3)  $\frac{25m}{24hR}$       (4)  $\frac{24m}{25hR}$

Scan or tap for the full solution



**Q.31** An electron in the hydrogen atom jumps from excited state  $n$  to the ground state. The wavelength so emitted illuminates a photosensitive material having work function 2.75 eV. If the stopping potential of the photoelectron is 10 V, then the value of  $n$  is

[AIPMT 2011 (Mains)]

- (1) 2      (2) 3      (3) 4      (4) 5

Scan or tap for the full solution



**Q.32** Out of the following which one is not a possible energy for a photon to be emitted by hydrogen atom according to Bohr's atomic model?

[AIPMT 2011 (Mains)]

- (1) 0.65 eV (2) 1.9 eV (3) 11.1 eV (4) 13.6 eV

Scan or tap for the full solution



**Q.33** The wavelength of the first line of Lyman series for hydrogen atom is equal to that of the second line of Balmer series for a hydrogen-like ion. The atomic number  $Z$  of hydrogen-like ion is

[AIPMT 2011 (Prelims)]

- (1) 3 (2) 4 (3) 1 (4) 2

Scan or tap for the full solution



**Q.34** The energy of a hydrogen atom in the ground state is  $-13.6$  eV. The energy of a  $\text{He}^+$  ion in the first excited state will be

[AIPMT 2010 (Prelims)]

- (1)  $-13.6$  eV (2)  $-27.2$  eV  
(3)  $-54.4$  eV (4)  $-6.8$  eV

Scan or tap for the full solution



**Q.35** The energy of a hydrogen atom in the ground state is  $-13.6$  eV. The energy of a  $\text{He}^+$  ion in the first excited state will be

[AIPMT 2010 (Prelims)]

- (1)  $-13.6$  eV (2)  $-27.2$  eV  
(3)  $-54.4$  eV (4)  $-6.8$  eV

Scan or tap for the full solution



**Q.36** An alpha nucleus of energy  $\frac{1}{2}mv^2$  bombards a heavy nuclear target of charge  $Ze$ . Then the distance of closest approach for the alpha nucleus will be proportional to

[AIPMT 2010 (Prelims)]

- (1)  $\frac{1}{Ze}$  (2)  $v^2$  (3)  $\frac{1}{m}$  (4)  $\frac{1}{v^4}$

Scan or tap for the full solution



**Q.37** In a Rutherford scattering experiment when a projectile of charge  $z_1$  and mass  $M_1$  approaches a target nucleus of charge  $z_2$  and mass  $M_2$ , the distance of closest approach is  $r_0$ . The energy of the projectile is

[AIPMT 2009]

- (1) directly proportional to  $z_1 z_2$   
(2) inversely proportional to  $z_1$   
(3) directly proportional to mass  $M_1$   
(4) directly proportional to  $M_1 \times M_2$

Scan or tap for the full solution



**Q.38** The ionization energy of the electron in the hydrogen atom in its ground state is  $13.6$  eV. The atoms are excited to higher energy levels to emit radiations of 6 wavelengths. Maximum wavelength of emitted radiation corresponds to the transition between

[AIPMT 2009]

- (1)  $n = 3$  to  $n = 1$  states (2)  $n = 2$  to  $n = 1$  states  
(3)  $n = 4$  to  $n = 3$  states (4)  $n = 3$  to  $n = 2$  states

Scan or tap for the full solution



**Q.39** In the phenomenon of electric discharge through gases at low pressure, the coloured glow in the tube appears as a result of

[AIPMT 2008]

- (1) collisions between the charged particles emitted from the cathode and the atoms of the gas  
(2) collision between different electrons of the atoms of the gas  
(3) excitation of electrons in the atoms  
(4) collision between the atoms of the gas

Scan or tap for the full solution



**Q.40** The ground state energy of hydrogen atom is  $-13.6$  eV. When its electron is in the first excited state, its excitation energy is

[AIPMT 2008]

- (1)  $10.2$  eV (2)  $0$  (3)  $3.4$  eV (4)  $6.8$  eV

Scan or tap for the full solution



**Q.41** The total energy of electron in the ground state of hydrogen atom is  $-13.6$  eV. The kinetic energy of an electron in the first excited state is

[AIPMT 2007]

- (1)  $6.8$  eV (2)  $13.6$  eV (3)  $1.7$  eV (4)  $3.4$  eV

Scan or tap for the full solution



**Q.42** Ionization potential of hydrogen atom is  $13.6$  eV. Hydrogen atoms in the ground state are excited by monochromatic radiation of photon energy  $12.1$  eV. According to Bohr's theory, the spectral lines emitted by hydrogen will be

[AIPMT 2006]

- (1) one (2) two (3) three (4) four

Scan or tap for the full solution



**Q.43** The total energy of an electron in the first excited state of hydrogen atom is about  $-3.4$  eV. Its kinetic energy in this state is

[AIPMT 2005]

- (1)  $3.4$  eV (2)  $6.8$  eV (3)  $-3.4$  eV (4)  $-6.8$  eV

Scan or tap for the full solution



- Q.44** Energy levels A, B and C of a certain atom corresponding to increasing values of energy i.e.  $E_A < E_B < E_C$ . If  $\lambda_1$ ,  $\lambda_2$  and  $\lambda_3$  are wavelengths of radiations corresponding to transitions C to B, B to A and C to A respectively, which of the following relations is correct? [AIPMT 2005]

(1)  $\lambda_3 = \lambda_1 + \lambda_2$       (2)  $\lambda_3 = \frac{\lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$   
 (3)  $\lambda_1 + \lambda_2 + \lambda_3 = 0$       (4)  $\lambda_3^2 = \lambda_1^2 + \lambda_2^2$

Scan or tap for the full solution



- Q.45** The Bohr model of atoms [AIPMT 2004]

- (1) Assumes that the angular momentum of electrons is quantized.  
 (2) Uses Einstein's photoelectric equation.  
 (3) Predicts continuous emission spectra for atoms.  
 (4) Predicts the same emission spectra for all types of atoms.

Scan or tap for the full solution



- Q.46** In which of the following systems will the radius of the first orbit ( $n = 1$ ) be minimum? [AIPMT 2003]

- (1) doubly ionized lithium  
 (2) singly ionized helium  
 (3) deuterium atom  
 (4) hydrogen atom

Scan or tap for the full solution



- Q.47** The energy of hydrogen atom in  $n^{\text{th}}$  orbit is  $E_n$ , then the energy in  $n^{\text{th}}$  orbit of singly ionised helium atom will be [AIPMT 2001]

(1)  $4E_n$       (2)  $E_n/4$       (3)  $2E_n$       (4)  $E_n/2$

Scan or tap for the full solution



- Q.48** Maximum frequency of emission is obtained for the transition [AIPMT 2000]

- (1)  $n = 2$  to  $n = 1$       (2)  $n = 6$  to  $n = 2$   
 (3)  $n = 1$  to  $n = 2$       (4)  $n = 2$  to  $n = 6$

Scan or tap for the full solution



- Q.49** When an electron makes a transition from  $n = 4$  to  $n = 2$ , then the emitted line spectrum will be [AIPMT 2000]

- (1) first line of Lyman series  
 (2) second line of Balmer series  
 (3) first line of Balmer series  
 (4) second line of Paschen series

Scan or tap for the full solution



## ANSWER KEY

| Ques. | 1  | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9  | 10 | 11 |
|-------|----|----|----|----|----|----|----|----|----|----|----|
| Ans.  | 2  | 1  | 4  | 1  | 4  | 3  | 2  | 2  | 4  | 1  | 3  |
| Ques. | 12 | 13 | 14 | 15 | 16 | 17 | 18 | 19 | 20 | 21 | 22 |
| Ans.  | 4  | 2  | 3  | 4  | 1  | 2  | 2  | 1  | 3  | 1  | 3  |
| Ques. | 23 | 24 | 25 | 26 | 27 | 28 | 29 | 30 | 31 | 32 | 33 |
| Ans.  | 2  | 4  | 3  | 3  | 4  | 4  | 4  | 1  | 3  | 3  | 4  |
| Ques. | 34 | 35 | 36 | 37 | 38 | 39 | 40 | 41 | 42 | 43 | 44 |
| Ans.  | 1  | 1  | 3  | 1  | 3  | 1  | 1  | 4  | 3  | 1  | 2  |
| Ques. | 45 | 46 | 47 | 48 | 49 |    |    |    |    |    |    |
| Ans.  | 1  | 1  | 1  | 1  | 2  |    |    |    |    |    |    |