

PYQSide

NEET LAWS OF MOTION — IMPULSE — PYQS

Max. Marks: 36

Questions: 9

Section A (MCQ)

- Q.1** A ball of mass 0.5 kg is dropped from a height of 40 m. The ball hits the ground and rises to a height of 10 m. The impulse imparted to the ball during its collision with the ground is (Take $g = 9.8 \text{ m/s}^2$)

[NEET (UG) 2025]

- (1) 0 (2) 84 N s (3) 21 N s (4) 7 N s

Scan or tap for the full solution



- Q.2** A ball of mass 0.15 kg is dropped from a height 10 m, strikes the ground and rebounds to the same height. The magnitude of impulse imparted to the ball is ($g = 10 \text{ m/s}^2$) nearly :

[NEET (UG) 2021]


- (1) 1.4 kg m/s (2) 0 kg m/s
(3) 4.2 kg m/s (4) 2.1 kg m/s

Scan or tap for the full solution



- Q.3** A rigid ball of mass m strikes a rigid wall at 60° and gets reflected without loss of speed as shown in the figure. The value of impulse imparted by the wall on the ball will be :

[NEET 2016 (Phase 2)]




- (1) mV (2) $2mV$ (3) $\frac{mV}{2}$ (4) $\frac{mV}{3}$

Scan or tap for the full solution



- Q.4** The force F acting on a particle of mass m is indicated by the force-time graph shown below. The change in momentum of the particle over the time interval from zero to 8 s is :

[AIPMT 2014]



- (1) 24 N s (2) 20 N s (3) 12 N s (4) 6 N s

Scan or tap for the full solution



- Q.5** A body of mass M hits normally a rigid wall with velocity V and bounces back with the same velocity. The impulse experienced by the body is

[AIPMT 2011 (Prelims)]

- (1) MV (2) $1.5MV$ (3) $2MV$ (4) zero

Scan or tap for the full solution



- Q.6** A 0.5 kg ball moving with a speed of 12 m/s strikes a hard wall at an angle of 30° with the wall. It is reflected with the same speed at the same angle. If the ball is in contact with the wall for 0.25 seconds, the average force acting on the wall is :

[AIPMT 2006]

- (1) 96 N (2) 48 N (3) 24 N (4) 12 N

Scan or tap for the full solution



- Q.7** An object of mass 3 kg is at rest. Now a force of $\vec{F} = 6t^2\hat{i} + 4t\hat{j}$ is applied on the object then velocity of object at $t = 3$ sec. is

[AIPMT 2002]

- (1) $18\hat{i} + 3\hat{j}$ (2) $18\hat{i} + 6\hat{j}$ (3) $3\hat{i} + 18\hat{j}$ (4) $18\hat{i} + 4\hat{j}$

Scan or tap for the full solution



- Q.8** A cricketer catches a ball of mass 150 g in 0.1 s moving with speed 20 m/s, then he experiences a force of

[AIPMT 2001]


- (1) 300 N (2) 30 N (3) 3 N (4) 0.3 N

Scan or tap for the full solution



- Q.9** A body of mass 3 kg hits a wall at an angle of 60° and returns at the same angle. The impact time was 0.2 s. The force exerted on the wall is

[AIPMT 2000]



(1) $150\sqrt{3} \text{ N}$

(2) $50\sqrt{3} \text{ N}$

(3) 100 N

(4) $75\sqrt{3} \text{ N}$

Scan or tap for the
full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9		
Ans.	3	3	1	3	3	3	2	2	1		