

PYQSide

NEET MOTION IN A PLANE — UNIFORM CIRCULAR MOTION (ANGULAR VELOCITY, CENTRIPETAL ACCELERATION) — PYQS

Max. Marks: 52

Questions: 13

Section A (MCQ)

Q.1 A particle moving with uniform speed in a circular path maintains: [NEET (UG) 2024]

- (1) Constant velocity
- (2) Constant acceleration
- (3) Constant velocity but varying acceleration
- (4) Varying velocity and varying acceleration

Scan or tap for the full solution



Q.2 Let ω_1 , ω_2 and ω_3 be the angular speed of the second hand, minute hand and hour hand of a smoothly running analog clock, respectively. If x_1 , x_2 and x_3 are their respective angular distances in 1 minute then the factor which remains constant (k) is [RE-NEET (UG) 2024]

- (1) $\frac{\omega_1}{x_1} = \frac{\omega_2}{x_2} = \frac{\omega_3}{x_3} = k$
- (2) $\omega_1 x_1 = \omega_2 x_2 = \omega_3 x_3 = k$
- (3) $\omega_1 x_1^2 = \omega_2 x_2^2 = \omega_3 x_3^2 = k$
- (4) $\omega_1^2 x_1 = \omega_2^2 x_2 = \omega_3^2 x_3 = k$

Scan or tap for the full solution



Q.3 Two particles A and B are moving in uniform circular motion in concentric circles of radii r_A and r_B with speed v_A and v_B respectively. Their time period of rotation is the same. The ratio of angular speed of A to that of B will be : [NEET (UG) 2019]

- (1) $v_A : v_B$
- (2) 1:1
- (3) $r_B : r_A$
- (4) $r_A : r_B$

Scan or tap for the full solution



Q.4 A particle moves so that its position vector is given by $\vec{r} = \cos \omega t \hat{x} + \sin \omega t \hat{y}$, where ω is a constant. Which of the following is true? [NEET 2016 (Phase 1)]

- (1) Velocity is perpendicular to $\vec{r}$ and acceleration is directed towards the origin.
- (2) Velocity is perpendicular to $\vec{r}$ and acceleration is directed away from the origin.
- (3) Velocity and acceleration both are perpendicular to $\vec{r}$
- (4) Velocity and acceleration both are parallel to $\vec{r}$

Scan or tap for the full solution




Q.5 A particle of mass 10 g moves along a circle of radius 6.4 cm with a constant tangential acceleration. What is the magnitude of this acceleration if the kinetic energy of the particle becomes equal to 8×10^{-4} J by the end of the second revolution after the beginning of the motion? [NEET 2016 (Phase 1)]

- (1) 0.18 m/s²
- (2) 0.2 m/s²
- (3) 0.1 m/s²
- (4) 0.15 m/s²

Scan or tap for the full solution



Q.6 In the given figure, $a = 15 \text{ m s}^{-2}$ represents the total acceleration of particle moving in the clockwise direction in a circle of radius $R = 2.5 \text{ m}$ at a given instant of time. The speed of the particle is [NEET 2016 (Phase 2)]



- (1) 4.5 m s⁻¹
- (2) 5.0 m s⁻¹
- (3) 5.7 m s⁻¹
- (4) 6.2 m s⁻¹

Scan or tap for the full solution



Q.7 The position vector of a particle $\vec{R}$ as a function of time is given by $\vec{R} = 4\sin(2\pi t)\hat{i} + 4\cos(2\pi t)\hat{j}$. Where R is in meters, t is in seconds and $\hat{i}$ and $\hat{j}$ denote unit vectors along x - and y -directions, respectively. Which one of the following statements is wrong for the motion of particle? [AIPMT 2015]

- (1) Magnitude of the velocity of particle is 8 meter/second.
 (2) Path of the particle is a circle of radius 4 meter.
 (3) Acceleration vector is along $-\vec{R}$.
 (4) Magnitude of acceleration vector is $\frac{v^2}{R}$ where v is the velocity of particle.

Scan or tap for the full solution



- Q.8** A particle moves in a circle of radius 5 cm with constant speed and time period 0.2π s. The acceleration of the particle is [AIPMT 2011 (Prelims)]
 (1) 15 m/s^2 (2) 25 m/s^2 (3) 36 m/s^2 (4) 5 m/s^2

Scan or tap for the full solution



- Q.9** A particle moves in x-y plane according to rule $x = a \sin \omega t$ and $y = a \cos \omega t$. The particle follows

[AIPMT 2010 (Mains)]

- (1) an elliptical path
 (2) a circular path
 (3) a parabolic path
 (4) a straight line path inclined equally to x and y-axes

Scan or tap for the full solution



- Q.10** A car runs at a constant speed on a circular track of radius 100 m, taking 62.8 seconds for every circular lap. The average velocity and average speed for each circular lap respectively is [AIPMT 2006]

- (1) 10 m/s, 0 (2) 0, 0
 (3) 0, 10 m/s (4) 10 m/s, 10 m/s.

Scan or tap for the full solution



- Q.11** A stone tied to the end of a string of 1 m long is whirled in a horizontal circle with a constant speed. If the stone makes 22 revolutions in 44 seconds, what is the magnitude and direction of acceleration of the stone? [AIPMT 2005]

- (1) $\pi^2 \text{ m s}^{-2}$
 and direction along the radius towards the centre
 (2) $\pi^2 \text{ m s}^{-2}$
 and direction along the radius away from the centre
 (3) $\pi^2 \text{ m s}^{-2}$
 and direction along the tangent to the circle
 (4) $\pi^2/4 \text{ m s}^{-2}$
 and direction along the radius towards the centre.

Scan or tap for the full solution



- Q.12** A particle moves along a circle of radius $\left(\frac{20}{\pi}\right) \text{ m}$ with constant tangential acceleration. If the velocity of the particle is 80 m/s at the end of the second revolution after motion has begun, the tangential acceleration is [AIPMT 2003]
 (1) 40 m/s^2 (2) $640\pi \text{ m/s}^2$
 (3) $160\pi \text{ m/s}^2$ (4) $40\pi \text{ m/s}^2$

Scan or tap for the full solution



- Q.13** Two particles having mass M and m are moving in a circular path having radius R and r . If their time periods are the same, then the ratio of their angular velocities will be [AIPMT 2001]

- (1) $\frac{r}{R}$ (2) $\frac{R}{r}$ (3) 1 (4) $\sqrt{\frac{R}{r}}$

Scan or tap for the full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9	10	11
Ans.	4	1	2	1	3	3	1	4	2	3	1
Ques.	12	13									
Ans.	1	3									