

PYQSide

NEET NUCLEI — MASS–ENERGY EQUIVALENCE AND BINDING ENERGY — PYQS

Max. Marks: 44

Questions: 11

Section A (MCQ)

Q.1 The energy equivalent of 0.5 g of a substance is :


[NEET (UG) 2020]

- (1) 4.5×10^{13} J (2) 1.5×10^{13} J
(3) 0.5×10^{13} J (4) 4.5×10^{16} J

Scan or tap for the
full solution



Q.2 The binding energy per nucleon of ${}^7_3\text{Li}$ and ${}^4_2\text{He}$ nuclei are 5.60 MeV and 7.06 MeV respectively. In the nuclear reaction



the value of energy Q released is [AIPMT 2014]

- (1) 19.6 MeV (2) -2.4 MeV
(3) 8.4 MeV (4) 17.3 MeV

Scan or tap for the
full solution



Q.3 The power obtained in a reactor using ${}^{235}\text{U}$ disintegration is 1000 kW. The mass decay of ${}^{235}\text{U}$ per hour is [AIPMT 2011 (Prelims)]

- (1) 10 microgram (2) 20 microgram
(3) 40 microgram (4) 1 microgram

Scan or tap for the
full solution



Q.4 The binding energy per nucleon in deuterium and helium nuclei are 1.1 MeV and 7.0 MeV, respectively. When two deuterium nuclei fuse to form a helium nucleus the energy released in the fusion is [AIPMT 2010 (Mains)]

- (1) 23.6 MeV (2) 2.2 MeV
(3) 28.0 MeV (4) 30.2 MeV

Scan or tap for the
full solution



Q.5 The mass of a ${}^7_3\text{Li}$ nucleus is 0.042 u less than the sum of the masses of all its nucleons. The binding energy per nucleon of ${}^7_3\text{Li}$ nucleus is nearly [AIPMT 2010 (Prelims)]

- (1) 46 MeV (2) 5.6 MeV
(3) 3.9 MeV (4) 23 MeV

Scan or tap for the
full solution



Q.6 If $M(A, Z)$, M_p and M_n denote the masses of the nucleus ${}^A_Z\text{X}$, proton and neutron respectively in units of u ($1 \text{ u} = 931.5 \text{ MeV}/c^2$) and BE represents its binding energy in MeV, then [AIPMT 2008]

- (1) $M(A, Z) = ZM_p + (A - Z)M_n - \text{BE}$
(2) $M(A, Z) = ZM_p + (A - Z)M_n + \text{BE}/c^2$
(3) $M(A, Z) = ZM_p + (A - Z)M_n - \text{BE}/c^2$
(4) $M(A, Z) = ZM_p + (A - Z)M_n + \text{BE}$

Scan or tap for the
full solution



Q.7 A nucleus ${}^A_Z\text{X}$ has mass represented by $M(A, Z)$. If M_p and M_n denote the mass of proton and neutron respectively and B.E. the binding energy in MeV, then [AIPMT 2007]

- (1) B.E. = $[ZM_p + (A - Z)M_n - M(A, Z)]c^2$
(2) B.E. = $[ZM_p + AM_p - M(A, Z)]c^2$
(3) B.E. = $M(A, Z) - ZM_p - (A - Z)M_n$
(4) B.E. = $[M(A, Z) - ZM_p - (A - Z)M_n]c^2$

Scan or tap for the
full solution



Q.8 The binding energy of deuteron is 2.2 MeV and that of ${}^4_2\text{He}$ is 28 MeV. If two deuterons are fused to form one ${}^4_2\text{He}$ then the energy released is [AIPMT 2006]

- (1) 30.2 MeV (2) 25.8 MeV
(3) 23.6 MeV (4) 19.2 MeV

Scan or tap for the
full solution



Q.9 If $M(A, Z)$, M_p and M_n denote the masses of the nucleus ${}^A_Z\text{X}$, proton and neutron respectively in units of u ($1 \text{ u} = 931.5 \text{ MeV}/c^2$) and BE represents its binding energy in MeV, then [AIPMT 2004]

- (1) $M(A, Z) = ZM_p + (A - Z)M_n - \text{BE}$
(2) $M(A, Z) = ZM_p + (A - Z)M_n + \text{BE}/c^2$
(3) $M(A, Z) = ZM_p + (A - Z)M_n - \text{BE}/c^2$
(4) $M(A, Z) = ZM_p + (A - Z)M_n + \text{BE}$

Scan or tap for the
full solution



Q.10 The mass of proton is 1.0073 u and that of neutron is 1.0087 u ($\text{u} = \text{atomic mass unit}$). The binding energy of ${}^4_2\text{He}$ is (Given helium nucleus mass $\approx 4.0015 \text{ u}$) [AIPMT 2003]

- (1) 0.0305 J (2) 0.0305 erg
 (3) 28.4 MeV (4) 0.061 u

Scan or tap for the
full solution



Q.11 M_n and M_p represent the mass of neutron and proton respectively. An element having mass M has N neutrons and Z protons, then the correct relation will be

[AIPMT 2001]

- (1) $M < \{N \cdot M_n + Z \cdot M_p\}$ (2) $M > \{N \cdot M_n + Z \cdot M_p\}$
 (3) $M = \{N \cdot M_n + Z \cdot M_p\}$ (4) $M = N\{M_n + M_p\}$

Scan or tap for the
full solution



ANSWER KEY

Ques.	1	2	3	4	5	6	7	8	9	10	11
Ans.	1	4	3	1	2	3	1	3	3	3	1